

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PUBLIC HEALTH REPORTS.

UNITED STATES.

Disinfection of vessels at Tampa Bay Quarantine on account of tuberculosis.

Asst. Surg. C. C. Pierce reports the disinfection of the Norwegian bark *Pallas* at the Tampa Bay Quarantine, Mullet Key, Fla., November 4, 1902, on account of a case of tuberculosis, and the master of the vessel was instructed as to precautions to prevent further infection. The vessel was fumigated on account of rats, 22 of which were collected and burned.

Disinfection of vessels at San Francisco on account of tubérculosis.

P. A. Surg. W. G. Stimpson reports the disinfection of 3 vessels during September, 1902, on account of the employment of tuberculous seamen on board.

Statistical reports of States and cities of the United States—Yearly and monthly.

Illinois—Springfield.—Month of October, 1902. Estimated population, 36,000. Total number of deaths, 50, including diphtheria, 1, and 7 from tuberculosis.

MASSACHUSETTS—Newton.—Month of October, 1902. Estimated population, 36,336. Total number of deaths, 35, including 4 from tuberculosis.

MICHIGAN.—Reports to the State board of health, Lansing, for the week ended November 1, 1902, from 69 observers, indicate that diarrhea, enteric fever, pneumonia, erysipelas, and measles were more prevalent and scarlet fever, inflammation of bowels, cholera morbus, and dysentery were less prevalent than in the preceding week. Meningitis was reported present at 3, measles at 18, smallpox at 19, whooping cough at 20, diphtheria at 43, scarlet fever at 63, enteric fever at 91, and phthisis pulmonalis at 226 places.

MINNESOTA—Winona.—Month of October, 1902. Estimated population, 23,000. Total number of deaths 24, including enteric fever, 1; whooping cough, 1, and 2 from tuberculosis.

NEW YORK.—Reports to the State board of health, Albany, for the month of September, 1902, from 156 cities, towns, and villages having an aggregate estimated population of 7,467,050, show a total of 9,682 deaths, including diphtheria, 166; enteric fever, 179; measles, 7; scarlet fever, 31; whooping cough, 83; smallpox, 6, and 917 from tuberculosis.

2595